

University of Groningen

The crystal structure of a hyperthermoactive exopolygalacturonase from *Thermotoga maritima* reveals a unique tetramer

Pijning, Tjaard; van Pouderoyen, Gertie; Kluskens, Leon; van der Oost, John; Dijkstra, Bauke W.

Published in:
FEBS Letters

DOI:
[10.1016/j.febslet.2009.10.047](https://doi.org/10.1016/j.febslet.2009.10.047)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Pijning, T., van Pouderoyen, G., Kluskens, L., van der Oost, J., & Dijkstra, B. W. (2009). The crystal structure of a hyperthermoactive exopolygalacturonase from *Thermotoga maritima* reveals a unique tetramer. *FEBS Letters*, 583(22), 3665-3670. <https://doi.org/10.1016/j.febslet.2009.10.047>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplementary data




Fig. S1. Sequence alignment of polygalacturonases, figure prepared by ESPrift. *Tma* = *Thermotoga maritima*, *Yen* = *Yersinia enterocolitica* (β-helical domain), *Eca* = *Erwinia carotovora*, *Ani* = *Aspergillus niger*, *Spu* = *Sterum purpureum*, *Clu* = *Colletotrichum lupine*, *Aac* = *Aspergillus aculeatus*, *Fmo* = *Fusarium moniliforme*. Fully conserved residues are boxed in red; the three catalytic aspartates are marked with ▲. Secondary structure elements are coded as follows: α = α-helix, β = β-strand, η = 310-helix, T = hydrogen bonded turn. Turns and loops lining the active site are indicated with their coil number and turn type.